## Recombinant Pasteurella haemolytica Leukotoxin(lktA)

Catalog No: #AP72878

Package Size: #AP72878-1 20ug #AP72878-2 100ug #AP72878-3 1mg



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

| Product Name          | Recombinant Pasteurella haemolytica Leukotoxin(lktA)   |
|-----------------------|--|
| Brief Description     | Recombinant Protein  |
| Host Species          | Yeast  |
| Purification          | Greater than 90% as determined by SDS-PAGE.  |
| Immunogen Description | Expression Region:715-953aaSequence Info:Partial   |
| Accession No.         | P0C085   |
| Uniprot               | P0C085   |
| Calculated MW         | 28 kDa   |
| Tag Info              | N-terminal 6xHis-tagged  |
| Target Sequence       | ${\tt IGTSHNDIFKGSKFNDAFNGGDGVDTIDGNDGNDRLFGGKGDDIIDGGNGDDFIDGGKGNDLLHGGKGDDIFV}$                                    |
|                       | HRQGDGNDSITESEGNDKLSFSDSNLKDLTFEKVNHHLVITNTKQEKVTIQNWFREAEFAKTIQNYVATRDDKI   |
|                       | EEIIGQNGERITSKQVDELIEKGNGKIAQSELTKVVDNYQLLKYSRDASNSLDKLISSASAFTSSNDSRNVLASP  |
|                       | TSMLDPSLSSIQFARAA  |
| Formulation           | Tris-based buffer50% glycerol  |
| Storage               | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability  |
|                       | of the protein itself.   |
|                       | Generally, the shelf life of liquid form is 6 months at -20°C,-80°C. The shelf life of lyophilized form is 12 months |
|                       | at -20°C,-80°C.Notes:Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for             |
|                       | up to one week.  |

## Background

Pasteurella leukotoxins are exotoxins that attack host leukocytes and especially polymorphonuclear cells, by causing cell rupture. The leukotoxin binds to the host LFA-1 integrin and induces a signaling cascade leading to many biological effects, including tyrosine phosphorylation of the CD18 tail, elevation of the intracellular Ca2+ and lysis of the host cell. This leukotoxin is a major contributor to the pathogenesis of lung injury in ovine pneumonic pasteurellosis. It has also weak holytic activity.

## References

Sequence diversity and molecular evolution of the leukotoxin (lktA) gene in bovine and ovine strains of Mannheimia (Pasteurella) haemolytica. Davies R.L., Whittam T.S., Selander R.K.J. Bacteriol. 183:1394-1404(2001) Research Topic: Others

Note: This product is for in vitro research use only