

Recombinant Influenza B virus Nucleoprotein

Catalog No: #AP72957



Package Size: #AP72957-1 20ug #AP72957-2 100ug #AP72957-3 1mg

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Recombinant Influenza B virus Nucleoprotein
Brief Description	Recombinant Protein
Host Species	Yeast
Purification	Greater than 90% as determined by SDS-PAGE.
Immunogen Description	Expression Region:1-168aaSequence Info:Partial
Other Names	Nucleocapsid protein ;Protein N
Accession No.	P04666
Uniprot	P04666
Calculated MW	20.8 kDa
Tag Info	N-terminal 6xHis-tagged
Target Sequence	MSNMDIDGINTGTIDKTPEEIISGTSGATRPIIRPATLAPPSNKRTRNPSPERATTSSEADVGRKTQKKQTPTTEIK KSVYNMVKLGEFYNQMMVKAGLNDDMERNLIQNAHAVERILLAATDDKKTEFQKKKNARDVKEGKEEIDHN KTGGTFYKMVRDDKTIYFSP
Formulation	Tris-based buffer50% glycerol
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C,-80°C. The shelf life of lyophilized form is 12 months at -20°C,-80°C.Notes:Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Background

Encapsidates the negative strand viral RNA, protecting it from nucleases. The encapsidated genomic RNA is termed the ribonucleoprotein (RNP) and serves as template for transcription and replication. The RNP needs to be localized in the nucleus to start an infectious cycle, but is too large to diffuse through the nuclear pore complex. NP comprises at least 2 nuclear localization signals and is responsible of the active RNP import into the nucleus through the cellular importin alpha,beta pathway. Later in the infection, nucleus export of RNP are mediated through viral proteins NEP interacting with M1 which binds nucleoproteins. It is possible that the nucleoprotein binds directly exportin-1 (XPO1) and plays an active role in RNP nuclear export. M1 interaction with RNP serves to hide nucleoprotein's nuclear localization signals. Soon after a virion infects a new cell, M1 dissociates from the RNP under acidification of the virion driven by M2 protein. Dissociation of M1 from RNP unmask nucleoprotein's nuclear localization signals, targeting the RNP to the nucleus .

References

Complete nucleotide sequence of the nucleoprotein gene of influenza B virus.Londo D.R., Davis A.R., Nayak D.P.J. Virol. 47:642-648(1983)Research Topic:Others

Note: This product is for in vitro research use only