

PDCD4 Antibody


Catalog No: #24468

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	PDCD4 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB IHC
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide
Immunogen Description	Raised against a 19 amino acid peptide from near the carboxy terminus of human PDCD4.
Target Name	PDCD4
Other Names	Programmed cell death 4, neoplastic transformation inhibitor
Accession No.	Swiss-Prot:Q53EL6Gene ID:27250
Uniprot	Q53EL6
GeneID	27250;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of PDCD4 in A-20 cell lysate with PDCD4 antibody at (A) 0.5 and (B) 1ug/mL.



Immunohistochemistry of PDCD4 in mouse brain tissue with PDCD antibody at 2.5 ug/mL.

Background

Apoptosis, also known as programmed cell death, plays major roles in development and normal tissue turnover in addition to tumor formation. During this process, the expression patterns of numerous genes are radically altered. One such gene is the programmed cell death protein 4 (PDCD4), whose expression was found to be upregulated in all cell lines following the onset of apoptosis. PDCD4 encodes a tumor suppressor protein whose expression is lost in carcinomas of breast, colon, lung and prostate. It can bind to and inhibit the helicase activity of the eukaryotic translation initiation factor 4A and inhibit the transactivation and transformation mediated by the transcription factor AP-1. The kinase Akt regulates PDCD4 by phosphorylation, decreasing the ability of PDCD4 to interfere with the transactivation of AP-1-responsive promoter by c-Jun. There are two known isoforms of PDCD4.

Note: This product is for in vitro research use only