JNK1/JNK2/JNK3(Ab-183/185) Antibody

Catalog No: #21504

Package Size: #21504-1 50ul #21504-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	JNK1/JNK2/JNK3(Ab-183/185) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	WB
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total JNK1/JNK2/JNK3(Ab-183/185) protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.182~186 (M-T-P-Y-V) derived from Human JNK1 JNK2 JNK3.
Target Name	JNK1 JNK2 JNK3
Other Names	PRKM8; SAPK1;PRKM9;SAPK1A;JNK3A
Accession No.	Swiss-Prot: P45983 P45984 P53779 NCBI Protein: NP_620637.1/NP_002743.3/NP_620448.1 NCBI
	mRNA:NM_139049.3/NM_002752.4/NM_138982.3
Uniprot	P45983
GeneID	5599;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.


Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 46/54kd
Western blotting: 1:500~1:1000

Images

Storage



Western blot analysis of extracts from 293 and Hela cells using JNK1/JNK2/JNK3(Ab-183/185) Antibody #21504.

Background

Serine/threonine-protein kinase involved in various processes such as cell proliferation, differentiation, migration, transformation and programmed cell death. Extracellular stimuli such as proinflammatory cytokines or physical stress stimulate the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway. In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK8/JNK1. In turn, MAPK8/JNK1 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity. Phosphorylates the replication licensing factor CDT1, inhibiting the interaction between CDT1 and the histone H4 acetylase HBO1 to replication origins. Loss of this interaction abrogates the acetylation required for replication initiation. Promotes stressed cell apoptosis by phosphorylating key regulatory factors including p53/TP53 and Yes-associates protein YAP1. In T-cells, MAPK8 and MAPK9 are required for polarized differentiation of T-helper cells into Th1 cells. Contributes to the survival of erythroid cells by phosphorylating the antagonist of cell death BAD upon EPO stimulation. Mediates starvation-induced BCL2 phosphorylation, BCL2 dissociation from BECN1, and thus activation of autophagy. Phosphorylates STMN2 and hence regulates microtubule dynamics, controlling neurite elongation in cortical neurons. In the developing brain, through its cytoplasmic activity on STMN2, negatively regulates the rate of exit from multipolar stage and of radial migration from the ventricular zone. Phosphorylates several other substrates including heat shock factor protein 4 (HSF4), the deacetylase SIRT1, ELK1, or the E3 ligase ITCH. Phosphorylates the CLOCK-ARNTL/BMAL1 heterodimer and plays a role in the regulation of the circadian clock (PubMed:22441692). Phosphorylates the heat shock transcription factor HSF1, suppressing HSF1-induced transcriptional activity (PubMed:10747973).

Note: This product is for in vitro research use only