

## Smad3(Phospho-Ser425) Antibody

Catalog No: #11325



Package Size: #11325-1 50ul #11325-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Smad3(Phospho-Ser425) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of Smad3 only when phosphorylated at serine 425.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 425 (C-S-S-V-S(p)) derived from Human Smad3.
Target Name	Smad3
Modification	Phospho
Other Names	JV15-2; MAD-3; MADH3; Mad3; Mothers against DPP homolog 3
Accession No.	Swiss-Prot: P84022NCBI Protein: NP_001138574.1
Uniprot	P84022
GeneID	4088;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


## Application Details

Predicted MW: 52kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

## Images



Western blot analysis of extracts from HeLa cells untreated or treated with EGF using Smad3(Phospho-Ser425) Antibody #11325.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Smad3(Phospho-Ser425) Antibody #11325(left) or the same antibody preincubated with blocking peptide(right).

## Background

Smad3 encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the *Drosophila* gene 'mothers against decapentaplegic' (Mad) and the *C. elegans* gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein functions as a transcriptional modulator activated by transforming growth factor-beta and is thought to play a role in the regulation of carcinogenesis.

Shi W, et al. *J Cell Sci.* 2007 Apr 1;120(Pt 7):1216-24

Seong HA, et al. *J Biol Chem.* 2007 Apr 20;282(16):12272-89

Wordinger RJ, et al. *Invest Ophthalmol Vis Sci.* 2007 Mar;48(3):1191-200

LeClair RJ, et al. *Circ Res.* 2007 Mar 30;100(6):826-33

Note: This product is for in vitro research use only