## SATB2 Rabbit mAb

Catalog No: #52055

Package Size: #52055-1 50ul #52055-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escri	ntı	Λn
$\boldsymbol{ u}$	COUL	рu	OH

Dan de et Norma	CATRO Relatives Ale	
Product Name	SATB2 Rabbit mAb	
Host Species	Recombinant Rabbit	
Clonality	Monoclonal antibody	
Clone No.	S02-1H5	
Isotype	Rabbit IgG	
Purification	Affinity Purified	
Applications	WB	
Species Reactivity	Human, Rat	
Immunogen Description	A synthetic peptide of human SATB2	
Conjugates	Unconjugated	
Modification	Unmodification	
Other Names	GLSS; SATB family member 2; SATB homeobox 2; SATB2; Special AT rich sequence binding protein 2;	
Accession No.	Swiss-Prot:Q9UPW6GeneID:23314	
Uniprot	Q9UPW6	
GeneID	23314	
Calculated MW	Calculated MW: 83 kDa; Observed MW: 83 kDa	
Concentration	0.3 mg/ml	
Formulation	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA	
orage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.		

## Application Details

WB: 1/2000

## Images



Western blot detection of SATB2 in Rat Brain lysates using SATB2 Rabbit mAb(1:1000 diluted). Predicted band size:83kDa. Observed band size:83kDa.

## Background

Swiss-Prot Acc.Q9UPW6. Binds to DNA, at nuclear matrix- or scaffold-associated regions. Thought to recognize the sugar-phosphate structure of double-stranded DNA. Transcription factor controlling nuclear gene expression, by binding to matrix attachment regions (MARs) of DNA and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin remodeling enzymes and also by recruiting corepressors (HDACs) or coactivators (HATs) directly to promoters and enhancers. Required for the initiation of the upper-layer neurons (UL1) specific genetic program and for the inactivation of deep-layer neurons (DL) and UL2 specific genes, probably by modulating BCL11B expression. Repressor of Ctip2 and regulatory determinant of corticocortical connections in the developing cerebral cortex. May play an important role in palate formation. Acts as a molecular node in a transcriptional network regulating skeletal development and osteoblast differentiation.

Note: This product is for in vitro research use only