HRH3 Rabbit mAb

Catalog No: #52345

Package Size: #52345-1 50ul #52345-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | HRH3 Rabbit mAb |
|-----------------------|--|
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal antibody |
| Clone No. | S04-3I4 |
| Isotype | Rabbit IgG |
| Purification | Affinity Purified |
| Applications | WB |
| Species Reactivity | Human,Mouse,Rat |
| Immunogen Description | A synthetic peptide of human HRH3/Histamine 3 Receptor |
| Conjugates | Unconjugated |
| Modification | Unmodification |
| Other Names | HH3R; GPCR97 |
| Accession No. | Swiss-Prot:Q9Y5N1GeneID:11255 |
| Uniprot | Q9Y5N1 |
| GeneID | 11255 |
| Calculated MW | Calculated MW: 49 kDa; Observed MW: 49 kDa |
| Concentration | 0.3 mg/ml |
| Formulation | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

Application Details

WB: 1/1000;

Images



Western blot detection of HRH3/Histamine 3 Receptor in K562,C6,3T3 cell lysates using HRH3/Histamine 3 Receptor Rabbit mAb(1:1000 diluted).Predicted band size:49kDa.Observed band size:49kDa.

Background

Swiss-Prot Acc.Q9Y5N1.The H3 subclass of histamine receptors could mediate the histamine signals in CNS and peripheral nervous system. Signals through the inhibition of adenylate cyclase and displays high constitutive activity (spontaneous activity in the absence of agonist). Agonist stimulation of isoform 3 neither modified adenylate cyclase activity nor induced intracellular calcium mobilization. Miscellaneous Does not bind to cimetidine and tripolidine. Shows modest affinity for thioperamide, imetit, N-alpha-methylhistamine and R--alpha-methylhistamine. Isoform 4 is unable to bind to iodoproxyfan while isoforms 1 and 3 bind it with high affinity.

Note: This product is for in vitro research use only