## NMDAR1 Rabbit mAb

Catalog No: #52464

Package Size: #52464-1 50ul #52464-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$				
	20	$\sim$ rı	m T I	n
	-	ULI	υu	on

Product Name	NMDAR1 Rabbit mAb	
Host Species	Recombinant Rabbit	
Clonality	Monoclonal antibody	
Clone No.	S08-4B6	
Isotype	Rabbit IgG	
Purification	Affinity Purified	
Applications	WB	
Species Reactivity	Human,Mouse,Rat	
Immunogen Description	A synthetic peptide of human NMDAR1	
Conjugates	Unconjugated	
Modification	Unmodification	
Other Names	NR1; MRD8; GluN1; NMDA1; NDHMSD; NDHMSR; NMD-R1; NMDAR1	
Accession No.	Swiss-Prot:Q05586GeneID:	
Uniprot	Q05586	
Calculated MW	Calculated MW: 105 kDa; Observed MW: 120 kDa	
Concentration	0.3 mg/ml	
Formulation	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.	

## **Application Details**

WB: 1/2000

## **Images**



Western blot detection of NMDAR1 in Rat brain, Jurkat cell lysates using NMDAR1 Rabbit mAb(1:1000 diluted). Predicted band size:105kDa. Observed band size:120kDa.

## Background

Swiss-Prot Acc.Q05586.Component of NMDA receptor complexes that function as heterotetrameric, ligand-gated ion channels with high calcium

permeability and voltage-dependent sensitivity to magnesium. Channel activation requires binding of the neurotransmitter glutamate to the epsilon subunit, glycine binding to the zeta subunit, plus membrane depolarization to eliminate channel inhibition by Mg2+ (PubMed:7685113, PubMed:28126851, PubMed:26919761, PubMed:26875626, PubMed:28105280). Sensitivity to glutamate and channel kinetics depend on the subunit composition (PubMed:26919761).

Note: This product is for in vitro research use only