V5-tag Rabbit mAb

Catalog No: #56317

Package Size: #56317-1 50ul #56317-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | | |
|------------------|------|-----|-----|-----|
| | esci | rın | tic | ۱n |
| \boldsymbol{L} | しつし | טוו | uv | , , |

| Product Name | V5-tag Rabbit mAb | |
|-----------------------|---|--|
| Host Species | Rabbit | |
| Clonality | Monoclonal | |
| Isotype | Rabbit IgG | |
| Purification | Affinity-chromatography | |
| Applications | WB | |
| Species Reactivity | All | |
| Specificity | V5 tag antibody detects endogenous levels of C-terminal, internal, and N-terminal V5-tagged proteins. | |
| Immunogen Description | A synthesized peptide | |
| Other Names | V5 tag | |
| Accession No. | Uniprot:N/A | |
| Uniprot | N/A | |
| Calculated MW | N/A | |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. | |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. | |
| | | |

Application Details

WB:1:3000~1:10000

Images



Western blot analysis of extracts from v5-tag fusion protein, using V5 tag antibody.

Product Description

The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids (GKPIPNPLLGLDST), although it has also been used with a shorter 9 amino acid sequence (IPNPLLGLD).

Note: This product is for in vitro research use only