

Histone H2B (Phospho-Thr129) Conjugated Antibody

Catalog No: #C14161



Package Size: #C14161-AF350 100ul #C14161-AF405 100ul #C14161-AF488 100ul
 #C14161-AF555 100ul #C14161-AF594 100ul #C14161-AF647 100ul
 #C14161-AF680 100ul #C14161-AF750 100ul #C14161-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	Histone H2B (Phospho-Thr129) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Species Reactivity	yeast
Specificity	Phospho-Histone H2B (T129) Antibody detects endogenous levels of total Phospho-Histone H2B (T129)
Immunogen Description	A synthesized peptide derived from yeast Phospho-Histone H2B (T129)
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	HTB2; Htb2p; HTB1; Htb1p; Histone H2B.1; Histone H2B.2; SPT12;
Accession No.	Uniprot:P02293(yeast)
Uniprot	P02293(yeast)
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	14kDa
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Note: This product is for in vitro research use only