

LRRK2 Conjugated Antibody

Catalog No: #C56018



Package Size: #C56018-AF350 100ul #C56018-AF405 100ul #C56018-AF488 100ul
 #C56018-AF555 100ul #C56018-AF594 100ul #C56018-AF647 100ul
 #C56018-AF680 100ul #C56018-AF750 100ul #C56018-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	LRRK2 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Species Reactivity	Human Mouse Rat
Specificity	LRRK2 Antibody detects endogenous levels of total LRRK2
Immunogen Description	A synthesized peptide derived from human LRRK2
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	AURA17; Dardarin antibody;; Leucine rich repeat kinase 2; LRRK 2 antibody; LRRK2; LRRK2_HUMAN; PARK 8; PARK8; RIPK7; ROCO 2; ROCO2;
Accession No.	Uniprot:Q5S007
Uniprot	Q5S007
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	280kDa
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250
 AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

Positively regulates autophagy through a calcium-dependent activation of the CaMKK/AMPK signaling pathway. The process involves activation of nicotinic acid adenine dinucleotide phosphate (NAADP) receptors, increase in lysosomal pH, and calcium release from lysosomes. Together with RAB29, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner.

Note: This product is for in vitro research use only