

Histone H3 (mono+di+tri methyl K14) Conjugated Antibody

Catalog No: #C56081

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C56081-AF350 100ul #C56081-AF405 100ul #C56081-AF488 100ul

#C56081-AF555 100ul #C56081-AF594 100ul #C56081-AF647 100ul

#C56081-AF680 100ul #C56081-AF750 100ul #C56081-Biotin 100ul

Description

Product Name	Histone H3 (mono+di+tri methyl K14) Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Species Reactivity	Human Mouse
Specificity	Histone H3 (mono+di+tri methyl K14) Antibody detects endogenous levels of total Histone H3 (mono+di+tri methyl K14)
Immunogen Description	A synthesized peptide derived from human Histone H3 (mono+di+tri methyl K14)
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	H3 histone; HIST1H3A; Histone cluster 1, H3a;H3/l; HIST3H3; H3K14me1; H3K14me2; H3K14me3;
Accession No.	Uniprot:P68431
Uniprot	P68431
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	15kDa
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Note: This product is for in vitro research use only