## LSD2 / AOF1 Conjugated Antibody

Catalog No: #C56262



 Package Size:
 #C56262-AF350 100ul
 #C56262-AF405 100ul
 #C56262-AF488 100ul

 #C56262-AF555 100ul
 #C56262-AF594 100ul
 #C56262-AF647 100ul

 #C56262-AF680 100ul
 #C56262-AF750 100ul
 #C56262-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

•	
Product Name	LSD2 / AOF1 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Species Reactivity	Human Mouse Rat
Specificity	LSD2 / AOF1 Antibody detects endogenous levels of total LSD2 / AOF1
Immunogen Description	A synthesized peptide derived from human LSD2 / AOF1
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	AOF1; KDM1B; LSD2; Lysine-specific histone demethylase 1B; Lysine-specific histone demethylase 2;
Accession No.	Uniprot:Q8NB78
Uniprot	Q8NB78
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	92kDa
Storage	Store at 4°C in dark for 6 months

## **Application Details**

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250		
AF405 conjugated: most applications: 1: 50 - 1: 250		
AF488 conjugated: most applications: 1: 50 - 1: 250		
AF555 conjugated: most applications: 1: 50 - 1: 250		
AF594 conjugated: most applications: 1: 50 - 1: 250		
AF647 conjugated: most applications: 1: 50 - 1: 250		
AF680 conjugated: most applications: 1: 50 - 1: 250		
AF750 conjugated: most applications: 1: 50 - 1: 250		
Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000		

## **Product Description**

Histone demethylase that demethylates 'Lys-4' of histone H3, a specific tag for epigenetic transcriptional activation, thereby acting as a corepressor. Required for de novo DNA methylation of a subset of imprinted genes during oogenesis. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed.

Note: This product is for in vitro research use only