Recombinant Rhesus Macaque Tumor Necrosis Factor-alpha/TNFSF2

SAB Signalway Antibody

Catalog No: #AP60148


Package Size: #AP60148-1 5ug #AP60148-2 100ug #AP60148-3 500ug

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|--------------|--|
| Draduat Nama | |

| Product Name | Recombinant Rhesus Macaque Tumor Necrosis Factor-alpha/TNFSF2 |
|-----------------|---|
| Host Species | Escherichia coli. |
| Purification | > 95 % by SDS-PAGE and HPLC analyses. |
| Other Names | TNFSF2, Cachectin, Differentiation-inducing Factor, DIF, Necrosin, Cytotoxin |
| Uniprot | P48094 |
| GeneID | 715467 |
| Calculated MW | Approximately 17.3 kDa, a single non-glycosylated polypeptide chain containing 157 amino acids. |
| Target Sequence | VRSSSRTPSD KPVAHVVANP QAEGQLQWLN RRANALLANG VELTDNQLVV PSEGLYLIYS |
| | QVLFKGQGCP SNHVLLTHTI SRIAVSYQTK VNLLSAIKSP CQRETPEGAE AKPWYEPIYL GGVFQLEKGD |
| | RLSAEINLPD YLDFAESGQV YFGIIAL |
| Formulation | Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4, 5 % trehalose. |
| Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles 12 months from date of receipt, -20 to |
| | -70 °C as supplied 1 month, 2 to 8 °C under sterile conditions after reconstitution 3 months, -20 to -70 °C |
| | under sterile conditions after reconstitution. |

Images



Background

Tumor necrosis factor alpha (TNF- α), also called cachectin, is the best-know member of the TNF-family, which can cause cell death. This protein is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF- α occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF- α is glycosylated, but non-glycosylated recombinant TNF- α has comparable biological activity. The biologically active native form of TNF- α is reportedly a trimer. Rhesus macaque and human TNF- α show approximately 98 % homology at the amino acid level. Two types of receptors for TNF- α have been described and virtually all cell types studied show the presence of one or both of these receptor types.

Note: This product is for in vitro research use only