TACC3 Rabbit mAb

Catalog No: #58603

Package Size: #58603-1 50ul #58603-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

Product Name	TACC3 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IHC ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	TACC3 Antibody detects endogenous levels of total TACC3
Immunogen Description	A synthesized peptide derived from human TACC3
Other Names	ERIC-1; ERIC1; TAC3; Transforming acidic coiled-coil-containing protein 3;TACC3;
Accession No.	Uniprot:Q9Y6A5
Uniprot	Q9Y6A5
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Application Details

WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

Images



Western blot analysis of TACC3 expression in C6 cell lysate.



Immunohistochemical analysis of paraffin-embedded human liver cancer, using TACC3 Antibody.

Product Description

TACC family members, TACC1, TACC2, and TACC3, map very closely to the corresponding FGFR1, FGFR2, FGFR3 genes on chromosomes 4,8, and 10. Subsequently, since they are phylogenetically related, it is proposed that TACC and FGFR have similar roles in cell growth and differentiation.

Background

TACC family members, TACC1, TACC2, and TACC3, map very closely to the corresponding FGFR1, FGFR2, FGFR3 genes on chromosomes 4,8, and 10. Subsequently, since they are phylogenetically related, it is proposed that TACC and FGFR have similar roles in cell growth and differentiation.

Note: This product is for in vitro research use only