BDNF Rabbit mAb

Catalog No: #58663

Signalway Antibody

Package Size: #58663-1 50ul #58663-2 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

Product Name	BDNF Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IHC ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	BDNF Antibody detects endogenous levels of total BDNF
Immunogen Description	A synthesized peptide derived from human BDNF
Other Names	BDNF;MGC34632;Abrineurin; ANON2; Brain Derived Neurotrophic Factor; Neurotrophin;BULN2;
Accession No.	Uniprot:P23560
Uniprot	P23560
Calculated MW	28kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Application Details

WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200


Images




Western blot analysis of extracts of Human cerebellum lysate, using BDNF antibody.



Immunohistochemical analysis of paraffin-embedded human testis, using BDNF Antibody .



Immunofluorescent analysis of HeLa cells, using BDNF Antibody.



All lanes use the Antibody at 1:1k dilution for 1 hour at room temperature.

Product Description

Neurotrophins function to regulate naturally occurring cell death of neurons during development. The prototype neurotrophin is nerve growth factor (NGF), originally discovered in the 1950s as a soluble peptide promoting the survival of, and neurite outgrowth from, sympathetic ganglia. Three additional structurally homologous neurotrophic factors have been identified. These include brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4) (also designated NT-5).

Background

Neurotrophins function to regulate naturally occurring cell death of neurons during development. The prototype neurotrophin is nerve growth factor (NGF), originally discovered in the 1950s as a soluble peptide promoting the survival of, and neurite outgrowth from, sympathetic ganglia. Three additional structurally homologous neurotrophic factors have been identified. These include brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin-4 (NT-4) (also designated NT-5).

Note: This product is for in vitro research use only