DKK1 Rabbit mAb

Catalog No: #58984

Package Size: #58984-1 50ul #58984-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


## Description

Product Name	DKK1 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IHC ICC/IF FC
Species Reactivity	Human Rat
Specificity	DKK1 Antibody detects endogenous levels of DKK1
Immunogen Description	A synthesized peptide derived from human DKK1
Other Names	SK; DKK-1; Dickkopf-related protein 1; Dickkopf-1; hDkk-1; Dickkopf homolog 1;
Accession No.	Uniprot:O94907
Uniprot	O94907
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


# Application Details

WB 1:1000~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50

### Images



Western blot analysis of DKK1 expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human placenta, using DKK1 Antibody.

### Product Description

DKK1: dickkopf homolog 1 (Xenopus laevis), also known as SK. Entrez Protein NP\_036374. DKK1 is a member of the dickkopf family. It is a secreted protein with two cysteine rich regions and is involved in embryonic development through its inhibition of the WNT signaling pathway. Elevated levels of DKK1 in bone marrow plasma and peripheral blood is associated with the presence of osteolytic bone lesions in patients with multiple myeloma.

#### Background

DKK1: dickkopf homolog 1 (Xenopus laevis), also known as SK. Entrez Protein NP\_036374. DKK1 is a member of the dickkopf family. It is a secreted protein with two cysteine rich regions and is involved in embryonic development through its inhibition of the WNT signaling pathway. Elevated levels of DKK1 in bone marrow plasma and peripheral blood is associated with the presence of osteolytic bone lesions in patients with multiple myeloma.

Note: This product is for in vitro research use only