FGFR3 Rabbit mAb

Catalog No: #59399

Package Size: #59399-1 50ul #59399-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_					
ш	es	cr	n	10	n
	-	C I I	IVI	II.	

Product Name	FGFR3 Rabbit mAb	
Host Species	Rabbit	
Clonality	Monoclonal	
Isotype	Rabbit IgG	
Purification	Affinity-chromatography	
Applications	WB	
Species Reactivity	Human Mouse Rat	
Specificity	FGFR3 Antibody detects endogenous levels of total FGFR3	
Immunogen Description	A synthesized peptide derived from human FGFR3	
Other Names	Fibroblast growth factor receptor 3; Heparin binding growth factor receptor; FGFR-3; CD333; FGFR3; JTK4;	
	FGFR 3; Tyrosine kinase JTK; $40\Omega\frac{1}{2}0\Omega\frac{1}{2}$	
Accession No.	Uniprot:P22607	
Uniprot	P22607	
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.	

Application Details

WB 1:500~1:2000

Images



Western blot analysis of FGFR3 expression in (1) A549 cell lysate; (2) Mouse brain lysate.

Product Description

FGFs act by binding and activating specific cell surface receptors. These include the Flg receptor or FGFR-1, the Bek receptor (or FGFR-2), FGFR-3, FGFR-4, FGFR-5 and FGFR-6. These receptors usually contain an extracellular ligandbinding region containing three immunoglobulin-like domains, a transmembrane domain and a cytoplasmic tyrosine kinase domain.

Background

FGFs act by binding and activating specific cell surface receptors. These include the FIg receptor or FGFR-1, the Bek receptor (or FGFR-2), FGFR-3, FGFR-4, FGFR-5 and FGFR-6. These receptors usually contain an extracellular ligandbinding region containing three immunoglobulin-like domains, a transmembrane domain and a cytoplasmic tyrosine kinase domain.

Note: This product is for in vitro research use only