

c-Rel Rabbit mAb

Catalog No: #59593

Package Size: #59593-1 50ul #59593-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	c-Rel Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IP
Species Reactivity	Human
Specificity	c-Rel Antibody detects endogenous levels of total c-Rel
Immunogen Description	A synthesized peptide derived from human c-Rel
Other Names	Avian reticuloendotheliosis; c Rel proto oncogene protein; Oncogene REL; Proto-oncogene c-Rel; REL;
Accession No.	Uniprot:Q04864
Uniprot	Q04864
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Application Details

WB 1:1000~1:5000 IP 1:50

Images



Western blot analysis of c-Rel expression in Daudi cell lysate.

Product Description

c-Rel contains an amino-terminal DNA-binding domain referred to as the REL homology domain (REH) and carboxy-terminal transactivation domains. The c-Rel protein is typically inhibited in unstimulated cells by I κ B α and I κ B β . c-Rel expression is highest in hematopoietic cells with extensive research studies demonstrating its role in immune cell function and pathogenesis of disease.

Background

c-Rel contains an amino-terminal DNA-binding domain referred to as the REL homology domain (REH) and carboxy-terminal transactivation domains. The c-Rel protein is typically inhibited in unstimulated cells by I κ B α and I κ B β . c-Rel expression is highest in hematopoietic cells with extensive research studies demonstrating its role in immune cell function and pathogenesis of disease.

Note: This product is for in vitro research use only