OGT Rabbit mAb

Catalog No: #59705

Package Size: #59705-1 50ul #59705-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_			
	Accr	iption	١
\boldsymbol{L}	COUL	เมนเบเ	ı

The state of the s		
Product Name	OGT Rabbit mAb	
Host Species	Rabbit	
Clonality	Monoclonal	
Isotype	Rabbit IgG	
Purification	Affinity-chromatography	
Applications	WB IHC ICC/IF FC	
Species Reactivity	Human Mouse Rat	
Specificity	OGT / O-Linked N-Acetylglucosamine Transferase Antibody detects endogenous levels of total OGT /	
	O-Linked N-Acetylglucosamine Transferase	
Immunogen Description	A synthesized peptide derived from human OGT / O-Linked N-Acetylglucosamine Transferase	
Other Names	HRNT1; O-GLCNAC;	
Accession No.	Uniprot:O15294	
Uniprot	O15294	
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.	

Application Details

WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50

Images



Western blot analysis of OGT / O-Linked N-Acetylglucosamine Transferase expression in A549 cell lysate.

Product Description

Addition of nucleotide-activated sugars directly onto the polypeptide through O-glycosidic linkage with the hydroxyl of serine or threonine. Mediates the O-glycosylation of MLL5 and HCFC1. Promotes proteolytic maturation of HCFC1.

Background

Addition of nucleotide-activated sugars directly onto the polypeptide through O-glycosidic linkage with the hydroxyl of serine or threonine. Mediates the O-glycosylation of MLL5 and HCFC1. Promotes proteolytic maturation of HCFC1.

Note: This product is for in vitro research use only