

Ube2N / Ubc13 Rabbit mAb

Catalog No: #59831

Package Size: #59831-1 50ul #59831-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Ube2N / Ubc13 Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB IHC ICC/IF
Species Reactivity	Human Mouse Rat
Specificity	Ube2N / Ubc13 Antibody detects endogenous levels of total Ube2N / Ubc13
Immunogen Description	A synthesized peptide derived from human Ube2N / Ubc13
Other Names	BLU; HEL-S-71; Ubc13; UbCH ben; UbCH13; UBCHBEN; Ube2n;
Accession No.	Uniprot:P61088
Uniprot	P61088
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Application Details

WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

Images



Western blot analysis of Ube2N / Ubc13 expression in Daudi cell lysate.

Product Description

The UBE2V1-UBE2N and UBE2V2-UBE2N heterodimers catalyze the synthesis of non-canonical 'Lys-63'-linked polyubiquitin chains. This type of polyubiquitination does not lead to protein degradation by the proteasome. Mediates transcriptional activation of target genes. Plays a role in the control of progress through the cell cycle and differentiation.

Background

The UBE2V1-UBE2N and UBE2V2-UBE2N heterodimers catalyze the synthesis of non-canonical 'Lys-63'-linked polyubiquitin chains. This type of polyubiquitination does not lead to protein degradation by the proteasome. Mediates transcriptional activation of target genes. Plays a role in the control of progress through the cell cycle and differentiation.

Note: This product is for in vitro research use only