UFD1L Rabbit mAb

Catalog No: #60188

Package Size: #60188-1 50ul #60188-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | UFD1L Rabbit mAb |
|----------------------|--|
| Host Species | Rabbit |
| Clonality | Monoclonal |
| sotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB ICC/IF FC |
| Species Reactivity | Human Mouse Rat |
| Specificity | UFD1L Antibody detects endogenous levels of total UFD1L |
| mmunogen Description | A synthesized peptide derived from human UFD1L |
| Other Names | UB fusion protein 1; UFD1; UFD1L; |
| Accession No. | Uniprot:Q92890 |
| Uniprot | Q92890 |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Application Details

WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:50

Images



Western blot analysis of UFD1L expression in HeLa cell lysate.

Product Description

Essential component of the ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. The ternary complex containing UFD1L, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome.

Background

Essential component of the ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. The ternary complex containing UFD1L, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome.

Note: This product is for in vitro research use only