

Bovine Fibrinogen Antibody FITC Conjugated

Catalog No: #C00716F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Bovine Fibrinogen Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IF
Species Reactivity	Cow
Immunogen Description	Fibrinogen from bovine plasma
Conjugates	FITC
Target Name	Bovine Fibrinogen
Other Names	FGA; FGA protein; FGB; FGG; Fib2; Fibrin alpha chain ;Fibrinogen A alpha polypeptide; Fibrinogen A alpha polypeptide chain; Fibrinogen alpha chain; Fibrinogen B alpha polypeptide; Fibrinogen beta chain; Fibrinogen G alpha polypeptide; Fibrinogen gamma chain; MGC104327; MGC119422; MGC119423; MGC11942
Accession No.	NCBI Gene ID522039
Uniprot	P02672
GeneID	522039;
Excitation Emission	494nm 518nm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

IF=1:50-200

Background

Fibrinogen is the main protein of blood coagulation system. It is a large protein and it consists of two identical subunits that contain three polypeptide chains: alpha, beta and gamma. All chains are connected with each other by a number of disulfide bonds. Fibrinopeptides A (1 to 16 amino acids) and B (1 to 17 amino acids) are released by thrombin from the N terminal parts of alpha and beta chains, respectively. In this way fibrinogen is converted into fibrin, which by means of polymerization forms a fibrin clot. Fibrinogen clotting underlies pathogenesis of MI, thromboembolism and thromboses of arteries and veins, since fibrin is the main substrate for thrombus formation. Fibrinogen activation is also involved in pathogenesis of inflammation, tumor growth and many other diseases. The normal fibrinogen concentration in plasma is about 3 mg/ml. The elevated level of fibrinogen in patient's blood is regarded as an independent risk factor for cardiovascular diseases. An increase in blood fibrinogen concentration was shown to be a strong predictor of coronary heart disease (Sonel A. et al, and Rapold H.J. et al). All these facts make fibrinogen an important parameter in the diagnosis of cardiovascular diseases.

Note: This product is for in vitro research use only