AMIGO1 Antibody FITC Conjugated

Catalog No: #C00928F



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech@signalwayantibody.com
Product Name	AMIGO1 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	ICC,IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human AMIGO1
Conjugates	FITC
Target Name	AMIGO1
Other Names	Adhesion molecule with Ig like domain 1; ali2; Alivin 2; AMIGO 1; AMIGO; AMIGO protein; Amphoterin
	induced gene and ORF Amigo; Amphoterin induced gene and ORF; Amphoterin induced protein 1; KIAA1163;
	AMGO1_HUMAN.
Excitation Emission	494nm 518nm
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

ICC=1:50-200 IF=1:50-200

Background

The amphoterin-induced gene and ORF (AMIGO) family of proteins consists of AMIGO-1, AMIGO-2 and AMIGO-3. All three members are single pass type I membrane proteins that contain several leucine-rich repeats, one IgG domain, and a transmembrane domain. The AMIGO proteins are specifically expressed on fiber tracts of neuronal tissues and participate in their formation. The AMIGO proteins can form complexes with each other, but can also bind itself. AMIGO-1, also designated Alivin-2, promotes growth and fasciculation of neurites and plays a role in myelination and fasciculation of developing neural axons. In cerebellar neurons, AMIGO-2 (Alivin-1) is crucial for depolarization-dependent survival. Similar to AMIGO-1 and AMIGO-2, AMIGO-3 (Alivin-3) plays a role in homophilic and or heterophilic cell-cell interaction and signal transduction.

Note: This product is for in vitro research use only