

Cdc25B(Phospho-Ser149) Antibody

Catalog No: #11553

Package Size: #11553-1 50ul #11553-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	Cdc25B(Phospho-Ser149) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB
Species Reactivity	Ms
Specificity	The antibody detects endogenous level of Cdc25B protein only when phosphorylated at serine 149.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of Serine 149 (F-R-S(p)-L-P) derived from Mouse Cdc25B.
Target Name	Cdc25B
Modification	Phospho
Other Names	CDC25B
Accession No.	Swiss-Prot:P30306Gene ID:12531
Uniprot	P30306
GeneID	12531;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 62kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from 3T3 cells untreated (lane 1) or treated with Hydroxyurea (lane 2) using Cdc25B(Phospho-Ser149) Antibody #11553.



Western blot analysis of extracts from HT29 cells, treated with PMA or calf intestinal phosphatase (CIP), using Cdc25B (Phospho-Ser149) Antibody #11553.

Background

Cdc25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined. Multiple transcript variants for this gene exist.

Li G, et al. (2010) *Retrovirology*.7:59.

Wang Z,et al. (2010) *BMC Cancer*.10:233.

Lucci MA,et al. (2010) *Cell Oncol*.32(5-6):361-72.

Note: This product is for in vitro research use only