PIRT Antibody FITC Conjugated

Catalog No: #C01356F



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | Support: tech@signalwayantibody.com |
|-----------------------|--|
| Product Name | PIRT Antibody FITC Conjugated |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | lgG |
| Purification | Purified by Protein A. |
| Applications | ICC IF |
| Species Reactivity | Hu Ms Rt |
| Immunogen Description | KLH conjugated synthetic peptide derived from human PIRT |
| Conjugates | FITC |
| Target Name | PIRT |
| Other Names | hCG_1776018; Phosphoinositide interacting regulator of transient receptor potential channels ; |
| | Phosphoinositide-interacting protein; Pirt; PIRT_HUMAN. |
| Excitation Emission | 494nm 518nm |
| Concentration | 1mg ml |
| Formulation | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |

Application Details

ICC=1:50-200 IF=1:50-200

Background

PIRT is a 137 amino acid multi-pass membrane protein. Highly conserved among vertebrates, PIRT consists of two transmembrane domains and one putative C-terminal phosphoinositide-binding domain. Although PIRT is expressed in peripheral nervous system, with highest levels in dorsal root ganglion and trigeminal neurons, and lowest levels in sympathetic and enteric neurons, it is not expressed in spinal cord. PIRT is a required component of the VR1 complex, which positively regulates VR1, a sensor of both noxious heat and capsaicin. Correspondingly, PIRT knockout results in impaired responses to noxious heat and capsaicin exposure, while VR1 remains unaltered. The gene that encodes PIRT maps to human chromosome 17p13.1.

Note: This product is for in vitro research use only