CD13 APN ANPEN Antibody PE Conjugated

Catalog No: #C02847P

Description



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|-----------------------|---|
| Product Name | CD13 APN ANPEN Antibody PE Conjugated |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Purified by Protein A. |
| Applications | Flow-Cyt IF |
| Species Reactivity | Hu Ms Rt |
| Immunogen Description | KLH conjugated synthetic peptide aa 910-960 967 derived from human CD13 |
| Conjugates | PE |
| Target Name | CD13 APN ANPEN |
| Other Names | APN; CD13; LAP1; P150; PEPN; GP150; Aminopeptidase N; AP-N; hAPN; Alanyl aminopeptidase; |
| | Aminopeptidase M; AP-M; Microsomal aminopeptidase; Myeloid plasma membrane glycoprotein CD13; |
| | ANPEP |
| Accession No. | Swiss-Prot#P15144NCBI Gene ID290 |
| Uniprot | P15144 |
| GeneID | 290; |
| Excitation Emission | 480,565nm 578nm |
| Cell Localization | Cytoplasm, Cell membrane |
| Concentration | 1mg ml |
| Formulation | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| | |

Application Details

Flow-Cyt=1:50-200 IF=1:50-200

Background

Broad specificity aminopeptidase. Plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. May play a critical role in the pathogenesis of cholesterol gallstone disease. May be involved in the metabolism of regulatory peptides of diverse cell types, responsible for the processing of peptide hormones, such as angiotensin III and IV, neuropeptides, and chemokines. Found to cleave antigen peptides bound to major histocompatibility complex class II molecules of presenting cells and to degrade neurotransmitters at synaptic junctions. Is also implicated as a regulator of IL-8 bioavailability in the endometrium, and therefore may contribute to the regulation of angiogenesis. Is used as a marker for acute myeloid leukemia and plays a role in tumor invasion. In case of human coronavirus 229E (HCoV-229E) infection, serves as receptor for HCoV-229E spike glycoprotein. Mediates as well human cytomegalovirus (HCMV) infection.

Note: This product is for in vitro research use only