

## CXCL10 Antibody FITC Conjugated

Catalog No: #C03000F

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

|                       |   |
|-----------------------|---|
| Product Name          | CXCL10 Antibody FITC Conjugated   |
| Host Species          | Rabbit  |
| Clonality             | Polyclonal  |
| Isotype               | IgG   |
| Purification          | Purified by Protein A.  |
| Applications          | IF  |
| Species Reactivity    | Hu Ms Rt  |
| Immunogen Description | KLH conjugated synthetic peptide aa 30-80 98 derived from mouse CXCL10  |
| Conjugates            | FITC  |
| Target Name           | CXCL10  |
| Other Names           | C7; IP1; CRG-2; INP1; IP-1; Ifi1; mob-1; Scyb1; gIP-1; C-X-C motif chemokine 1; 1 kDa interferon gamma-induced protein; Gamma-IP1; Interferon-gamma induced protein CRG-2; Small-inducible cytokine B1; Cxcl1; Crg2 |
| Accession No.         | Swiss-Prot#P17515NCBI Gene ID15945  |
| Uniprot               | P17515  |
| GeneID                | 15945;  |
| Excitation Emission   | 494nm 518nm   |
| Cell Localization     | Secreted  |
| Concentration         | 1mg ml  |
| Formulation           | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.  |
| Storage               | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.   |

## Application Details

IF=1:50-200

## Background

In addition to its role as a proinflammatory cytokine, may participate in T-cell effector function and perhaps T-cell development.

Note: This product is for in vitro research use only