

## VEGFR3 Antibody FITC Conjugated

Catalog No: #C03998F

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

Product Name	VEGFR3 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 910-960 1298 derived from human VEGFR-3
Conjugates	FITC
Target Name	VEGFR3
Other Names	PCL; FLT41; LMPH1A; VEGFR3; Vascular endothelial growth factor receptor 3; VEGFR-3; Fms-like tyrosine kinase 4; FLT-4; Tyrosine-protein kinase receptor FLT4; FLT4
Accession No.	Swiss-Prot#P35916NCBI Gene ID2324
Uniprot	P35916
GeneID	2324;
Excitation Emission	494nm 518nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## Application Details

Flow-Cyt=1:50-200 ICC=1:50-200 IF=1:50-200

## Background

Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFC and VEGFD, and plays an essential role in adult lymphangiogenesis and in the development of the vascular network and the cardiovascular system during embryonic development. Promotes proliferation, survival and migration of endothelial cells, and regulates angiogenic sprouting. Signaling by activated FLT4 leads to enhanced production of VEGFC, and to a lesser degree VEGFA, thereby creating a positive feedback loop that enhances FLT4 signaling. Modulates KDR signaling by forming heterodimers. The secreted isoform 3 may function as a decoy receptor for VEGFC and or VEGFD and play an important role as a negative regulator of VEGFC-mediated lymphangiogenesis and angiogenesis. Binding of vascular growth factors to isoform 1 or isoform 2 leads to the activation of several signaling cascades; isoform 2 seems to be less efficient in signal transduction, because it has a truncated C-terminus and therefore lacks several phosphorylation sites. Mediates activation of the MAPK1 ERK2, MAPK3 ERK1 signaling pathway, of MAPK8 and the JUN signaling pathway, and of the AKT1 signaling pathway. Phosphorylates SHC1. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Promotes phosphorylation of MAPK8 at 'Thr-183' and 'Tyr-185', and of AKT1 at 'Ser-473'.

---

Note: This product is for in vitro research use only