

p38 MAPK (Phospho-Thr180 + Tyr182) Antibody HRP Conjugated



Catalog No: #C04000H

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	p38 MAPK (Phospho-Thr180 + Tyr182) Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WB IHC-P IHC-F
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic phosphopeptide aa 160-200 360 derived from human p38 MAPK around the phosphorylation site of(Thr180 Tyr182)
Conjugates	HRP
Target Name	p38 MAPK Thr180 + Tyr182
Other Names	RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA; Mitogen-activated protein kinase 14; MAP kinase 14; MAPK 14; Cytokine suppressive anti-inflammatory drug-binding protein; CSAID-binding protein; MAP kinase MXI2; MAX-interacting protein 2; Mitogen-activated protein kin
Accession No.	Swiss-Prot#Q16539NCBI Gene ID1432
Uniprot	Q16539
GeneID	1432;
Excitation Emission	N A
Cell Localization	Cytoplasm, Nucleus
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

WB=1:500-2000 IHC-P=1:50-200 IHC-F=1:50-200

Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1 TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Note: This product is for in vitro research use only