CCL17 ANTIBODY FITC Conjugated

Catalog No: #C04117F



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech@signalwayantibody.com
Product Name	CCL17 ANTIBODY FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 40-90 103 derived from mouse CCL17
Conjugates	FITC
Target Name	CCL17
Other Names	Tarc; Abcd-2; Scya17; Scya17I
Accession No.	NCBI Gene ID20295
GeneID	20295;
Excitation Emission	494nm 518nm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

IF=1:50-200

Background

Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. TARC displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. TARC binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

Note: This product is for in vitro research use only