JAK2(Phospho-Tyr1007+Tyr1008) Antibody HRP Conjugated

NCBI Gene ID3717

O60674

3717;

1mg ml

NΑ

SAB Signalway Antibody

Catalog No: #C04135H

Description

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	JAK2(Phospho-Tyr1007+Tyr1008) Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WB IHC-P IHC-F
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic phosphopeptide derived from human JAK2 around the phosphorylation site of
	Tyr1007 1008
Conjugates	HRP
Target Name	JAK2 Tyr1007+Tyr1008
Other Names	JAK2Tyr1007 1008; Tyrosine protein kinase JAK2; JAK 2; JAK-2; JAK2; JAK2_HUMAN; Janus Activating
	Kinase 2; Janus Kinase 2; JTK 10; JTK10; OTTHUMP00000043260; Tyrosine-protein kinase JAK2; Tyrosine
	protein kinase JAK2.

Application Details

Accession No.

Excitation Emission

Concentration

Formulation

Storage

Uniprot GeneID

WB=1:500-2000 IHC-P=1:50-200 IHC-F=1:50-200

Background

JAK2 (Janus Activating Kinase 2) is a tyrosine kinase of the non-receptor type, that associates with the intracellular domains of cytokine receptors; JAK2 is the predominant JAK kinase activated in response to several growth factors and cytokines such as IL-3, GM-CSF and erythropoietin; it has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon. Ligand binding to a variety of cell surface receptors (e.g., cytokine, growth factor, GPCRs) leads to an association of those receptors with JAK proteins, which are then activated via phosphorylation on tyrosines 1007 and 1008 in the kinase activation loop. Activated JAK proteins phosphorylate and activate STAT (signal transducers and activators of transcription) proteins, which then dimerize and translocate to the nucleus. Once in the nucleus, STAT proteins bind to DNA and modify the transcription of various genes.

0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Note: This product is for in vitro research use only