

CK10 Antibody FITC Conjugated

Catalog No: #C04275F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	CK10 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt IF
Species Reactivity	HuB MsB RtB B
Immunogen Description	KLH conjugated synthetic peptide aa 140-190 584 derived from human CK10
Conjugates	FITC
Target Name	CK10
Other Names	BIE; EHK; K10; KPP; BCIE; CK10; Keratin, type I cytoskeletal 10; Cytokeratin-10; CK-10; Keratin-10; KRT10
Accession No.	Swiss-Prot#P13645NCBI Gene ID3858
Uniprot	P13645
GeneID	3858;
Excitation Emission	494nm 518nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1:50-200B IF=1:50-200B

Background

Cytokeratin 10 is a heterotetramer of two type I and two type II keratins. Cytokeratin 10 is generally associated with keratin 1. It is seen in all suprabasal cell layers including stratum corneum. A number of alleles are known that mainly differ in the Gly-rich region (positions 490-560). Defects in cytokeratin 10 are a cause of epidermolytic hyperkeratosis (EHK), also known as bullous congenital ichthyosiform erythroderma (BCIE) or bullous erythroderma ichthyosiformis congenita of Brocq. EHK is an hereditary skin disorder characterized by blistering and a marked thickening of the stratum corneum. At birth, affected individuals usually present with redness, blisters and superficial erosions due to cytolysis. Within a few weeks, the erythroderma and blister formation diminish and hyperkeratoses develop. Transmission is autosomal dominant, but most cases are sporadic. Defects in cytokeratin 10 are also a cause of annular epidermolytic ichthyosis (AEI), also known as cyclic ichthyosis with epidermolytic hyperkeratosis. AEI resembles clinical and histologic features of both epidermolytic hyperkeratosis and ichthyosis bullosa of Siemens.

Note: This product is for in vitro research use only