## CREB(Phospho-Ser142) Antibody

Catalog No: #11300

Package Size: #11300-1 50ul #11300-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description           |  |
|-----------------------|--|
| Product Name          | CREB(Phospho-Ser142) Antibody  |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.       |
|                       | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
|                       | specific antibodies were removed by chromatogramphy using non-phosphopeptide.                          |
| Applications          | WB IHC   |
| Species Reactivity    | Hu Ms Rt   |
| Specificity           | The antibody detects endogenous level of CREB only when phosphorylated at serine142.                   |
| Immunogen Type        | Peptide-KLH  |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 142(D-L-S(p)-S-D) derived from Human CREB.      |
| Target Name           | CREB   |
| Modification          | Phospho  |
| Other Names           | CREB   |
| Accession No.         | Swiss-Prot: P16220; NCBI Gene ID: 1385; NCBI mRNA: NM_004379.3; NCBI Protein: NP_004370.1              |
| Uniprot               | P16220   |
| GeneID                | 1385;  |
| SDS-PAGE MW           | 43KD   |
| Concentration         | 1.0mg/ml   |
| Formulation           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%   |

sodium azide and 50% glycerol.

Store at -20°C


## **Application Details**

Predicted MW: 43kd


Immunohistochemistry: 1:50~1:100

## **Images**

Storage



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using CREB (Phospho-Ser142) Antibody (#11300).



Western blot analysis of extracts from 293 cells, treated with calf intestinal phosphatase (CIP), using CREB(Phospho-Ser142) Antibody #11300.

## Background

Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells.

Note: This product is for in vitro research use only