

TK (Phospho-Ser13) Antibody

Catalog No: #12027

Package Size: #12027-1 50ul #12027-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	TK (Phospho-Ser13) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of TK only when phosphorylated at Serine 13.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of Serine 13(P-G-S(p)-P-S) derived from Human TK.
Target Name	TK
Modification	Phospho
Other Names	Thymidine kinase, cytosolic; TK1
Accession No.	Swiss-Prot#: P04183; NCBI Gene#: 7083; NCBI Protein#: XP_005257688.1
Uniprot	P04183
GeneID	7083;
SDS-PAGE MW	25kd
Concentration	1.0mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C/1 year

Application Details


Western blotting: 1:500~1:2000

Immunohistochemistry: 1:100~1:300

Images



Western blot analysis of lysates from HeLa cells treated with paclitaxel 1uM 24h, using TK (Phospho-Ser13) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using TK (Phospho-Ser13) Antibody. The picture on the right is blocked with the phospho peptide.

Background

TK is a cytosolic thymidine kinase. Phosphorylated during mitosis. Its enzymatic activity is high in proliferating cells and peaks during the S-phase of the cell cycle; it is very low in resting cells.

Note: This product is for in vitro research use only