GR (Phospho-Ser203) Antibody

Catalog No: #12062

Package Size: #12062-1 50ul #12062-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	GR (Phospho-Ser203) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of GR only when phosphorylated at Serine 203.
mmunogen Type	Peptide-KLH
mmunogen Description	Peptide sequence around phosphorylation site of Serine 203
	(S-G-S(p)-P-G) derived from Human GR.
Target Name	GR
Modification	Phospho
Other Names	GCCR, GCR, GR, GRL
Accession No.	Swiss-Prot#: P04150; NCBI Gene#: 2908; NCBI Protein#: NP_000167.1
Jniprot	P04150
GeneID	2908;
SDS-PAGE MW	86kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.029
	sodium azide and 50% glycerol.

Application Details


Predicted MW: 86kd

Western blotting: 1:500~1:1000

Images

Storage

Store at -20°C/1 year



Western blot analysis of extracts from A549 cells using GR (Phospho-Ser203) Antibody #12062. The lane on the left is treated with the antigen-specific peptide.

Background

Receptor for glucocorticoids (GC). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE), both for nuclear and mitochondrial DNA, and as a modulator of other transcription factors. Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth. Involved in chromatin remodeling. Plays a significant role in transactivation.

Note: This product is for in vitro research use only