## Histone H4K20me1 Polyclonal Antibody

Catalog No: #HW031

Package Size: #HW031-1 50ul #HW031-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_				
	escr	ın.	tio	m
$\boldsymbol{L}$	COUL	ıv	เเบ	48

Product Name	Histone H4K20me1 Polyclonal Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Purification	Affinity purification	
Applications	WB,IHC,IF	
Species Reactivity	Human,Mouse,Rat	
Immunogen Type	Peptide	
Immunogen Description	A synthetic peptide of human MonoMethyl-Histone H4-K20	
Target Name	Histone H4	
Modification	Methyl	
Other Names	FO108;H4;H4/n;H4F2;H4FN;HIST2H4;Histone H4;HIST1H4A;HIST2H4A	
Accession No.	Uniprot:P62805GeneID:8370	
Uniprot	P62805	
GeneID	8370	
SDS-PAGE MW	11KDa	
Concentration	1.0mg/ml	
Formulation	PBS with 0.02% sodium azide,50% glycerol,pH7.3.	
Storage	Store at -20°C. Avoid freeze / thaw cycles.	


## Application Details

WB 1:500 - 1:2000IHC 1:50 - 1:200IF 1:50 - 1:200


## **Images**




Western blot analysis of extracts of various cell lines, using MonoMethyl-Histone H4-K20 antibody.




Immunohistochemistry of paraffin-embedded rat testis using MonoMethyl-Histone H4-K20 antibody.




Immunofluorescence analysis of 293T cells using MonoMethyl-Histone H4-K20 antibody.



Immunohistochemistry of paraffin-embedded human colon using MonoMethyl-Histone H4-K20 antibody.



Immunohistochemistry of paraffin-embedded mouse testis using MonoMethyl-Histone H4-K20 antibody.



Western blot analysis of extracts of various cell lines, using MonoMethyl-Histone H4-K20 antibody.

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.

Note: This product is for in vitro research use only