AMPK beta 1 Monoclonal Antibody

Catalog No: #27201

Package Size: #27201-1 50ul #27201-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| Description | |
|-----------------------|--|
| Product Name | AMPK beta 1 Monoclonal Antibody |
| Host Species | Mouse |
| Clonality | Monoclonal |
| Clone No. | 1A7-E11-E9 |
| Isotype | Hu Ms Rt Mk |
| Purification | Affinity purified |
| Applications | WB ICC IP IHC |
| Species Reactivity | Hu Ms Rt Mk |
| Specificity | This antibody detects endogenous levels of AMPK beta 1 and does not cross-react with related proteins. |
| Immunogen Type | Recombinant Protein |
| Immunogen Description | Purified recombinant human AMPK beta 1 protein fragments expressed in E.coli. |
| Target Name | AMPK beta 1 |
| Other Names | 1300015D22Rik; 5 AMP activated protein kinase subunit beta 1; 5"-AMP-activated protein kinase subunit |
| | beta-1; AAKB1_HUMAN; AMP-ACTIVATED PROTEIN KINASE, NONCATALYTIC, BETA-1; AMP-activated, |
| | noncatalytic, beta-1; AMPK; AMPK beta 1 chain; |
| Accession No. | Uniprot: Q9Y478 Gene ID: 5564 |
| Uniprot | Q9Y478 |
| GenelD | 5564; |
| SDS-PAGE MW | 38kd |
| Formulation | Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.02% sodium azide and 50% glycerol. |
| Storage | store at -20A C |

Application Details


Western blotting: 1:1000

Immunocytochemistry: 1:100


Images




Immunocytochemistry staining of HeLa cells fixed with 1% Paraformaldehyde and using anti-AMPK beta 1 antibody (dilution 1:100).



Western blot detection of AMPK beta 1 in 3T3,Hela,PC-12,COS7 and MDA-MB-468 cell lysates using AMPK beta 1 antibody (1:1000 diluted).Predicted band size:38KDa.Observed band size:38KDa.Exposure time:5min.



Immunoprecipitation analysis of Hela cell lysates using AMPK beta 1 antibody.



Immunohistochemical analysis of paraffin-embedded Breast cancer using AMPK beta 1 Mouse mAb (1/200 dilution).Antigen retrieval was performed by pressure cooking in citrate buffer (pH 6.0).

Background

Non-catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Beta non-catalytic subunit acts as a scaffold on which the AMPK complex assembles, via its C-terminus that bridges alpha (PRKAA1 or PRKAA2) and gamma subunits (PRKAG1, PRKAG2 or PRKAG3).

Note: This product is for in vitro research use only