

DDB1 Monoclonal Antibody

Catalog No: #27205

Package Size: #27205-1 50ul #27205-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	DDB1 Monoclonal Antibody
Host Species	Mouse
Clonality	Monoclonal
Clone No.	2D6-B5-A8
Isotype	IgG2b
Applications	WB
Species Reactivity	Hu Ms Mk Rt
Specificity	This antibody detects endogenous levels of DDB1 and does not cross-react with related proteins.
Immunogen Type	Recombinant Protein
Immunogen Description	Purified recombinant human DDB1 protein fragments expressed in E.coli.
Target Name	DDB1
Other Names	Damage specific DNA binding protein 1; Damage-specific DNA-binding protein 1; DDB 1; DDB p127 subunit; Ddb1; DDB1_HUMAN; DDBa; DNA damage binding protein 1; DNA damage-binding protein 1; DNA damage-binding protein a; HBV X-associated protein 1;
Accession No.	Uniprot: Q16531 Gene ID: 1642
Uniprot	Q16531
GeneID	1642;
SDS-PAGE MW	127kd
Formulation	ascites
Storage	store at -20Λ C

Application Details

Western blotting: 1:1000

Images



Western blot detection of DDB1 antibody in HeLa,3T3,C6,COS7,293T and K562 cell lysates using DDB1 antibody (1:1000 diluted).Predicted band size:127KDa.Observed band size:127KDa.

Background

Required for DNA repair. Binds to DDB2 to form the UV-damaged DNA-binding protein complex (the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair. The UV-DDB complex preferentially binds to cyclobutane pyrimidine dimers (CPD), 6-4 photoproducts (6-4 PP), apurinic sites and short mismatches. Also appears to function as a component of numerous distinct DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. The functional specificity of the DCX E3 ubiquitin-protein ligase complex is determined by the variable substrate recognition component recruited by DDB1. DCX(DDB2) (also known as DDB1-CUL4-ROC1, CUL4-DDB-ROC1 and CUL4-DDB-RBX1) may ubiquitinate histone H2A, histone H3 and histone H4 at sites of UV-induced DNA damage. The ubiquitination of histones may facilitate their removal from the nucleosome and promote subsequent DNA repair. DCX(DDB2) also ubiquitinates XPC, which may enhance DNA-binding by XPC and promote NER. DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of TP53 in response to radiation-induced DNA damage and during DNA replication. DCX(ERCC8) (the CSA complex) plays a role in transcription-coupled repair (TCR). May also play a role in ubiquitination of CDKN1B/p27kip when associated with CUL4 and SKP2.

Note: This product is for in vitro research use only