

NMDAR1 (Phospho-Ser890) Antibody

Catalog No: #11674



Package Size: #11674-1 50ul #11674-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


| | |
|-----------------------|--|
| Product Name | NMDAR1 (Phospho-Ser890) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide. |
| Applications | IHC IF |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous levels of NMDAR1 only when phosphorylated at serine 890. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of Serine 890(A-S-S(p)-F-K) derived from Human NMDAR1. |
| Target Name | NMDAR1 |
| Modification | Phospho |
| Other Names | GLURZ1; GRIN1; NMD-R1; NMDZ1; NMZ1 |
| Accession No. | Swiss-Prot#: Q05586; NCBI Gene#: 2902; NCBI Protein#: NP_015566.1. |
| Uniprot | Q05586 |
| GeneID | 2902; |
| SDS-PAGE MW | 120kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at -20°C/1 year |

Application Details


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200

Images



Immunohistochemical analysis of paraffin-embedded human brain tissue using NMDAR1 (Phospho-Ser890) antibody #11674 (left) or the same antibody preincubated with blocking peptide (right).



Immunofluorescence staining of methanol-fixed A549 cells using NMDAR1 (Phospho-Ser890) Antibody #11674.

Background

NMDA receptors are members of the ionotropic class of glutamate receptors, which also includes Kainate and AMPA receptors. NMDA receptors consist of NR1 subunits combined with one or more NR2 (A-D) or NR3 (A-B) subunits. The ligand-gated channel is permeable to cations including Ca^{2+} , and at resting membrane potentials NMDA receptors are inactive due to a voltage-dependent blockade of the channel pore by Mg^{2+} . NMDA receptor activation, which requires binding of glutamate and glycine, leads to an influx of Ca^{2+} into the postsynaptic region where it activates several signaling cascades, including pathways leading to the induction of long-term potentiation (LTP) and depression (LTD). NMDA receptors have a critical role in excitatory synaptic transmission and plasticity in the CNS. They govern a range of physiological conditions including neurological disorders caused by excitotoxic neuronal injury, psychiatric disorders and neuropathic pain syndromes.

Guang Bai, J. Biol. Chem., Jan 1998; 273: 1086.

Guang Bai, J. Biol. Chem., Mar 1995; 270: 7737.

Mai T. Dang, PNAS, Oct 2006; 103: 15254 - 15259.

Note: This product is for in vitro research use only