## Sumo 2+3 Antibody HRP Conjugated

Catalog No: #C06832H



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech e signalwayantibouy.com
Product Name	Sumo 2+3 Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WB IHC-P IHC-F ICC
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human Sumo 2+3
Conjugates	HRP
Target Name	Sumo 2+3
Other Names	HSMT3; MGC117191; OTTHUMP00000115275; OTTHUMP00000115276; OTTHUMP00000115277; Sentrin
	2; Small ubiquitin like modier 2; Small ubiquitin like modier protein 3; Small ubiquitin related modier 2; Small
	ubiquitin related modier 3; SMT3 A; SMT3 B; SMT3 H1; SMT3 H2; SMT3 homolog 1; SMT3 homolog 2; SMT3
	ho
Excitation Emission	N A
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

WB=1:500-2000 IHC-P=1:50-200 IHC-F=1:50-200 ICC=1:50-200

## Background

The small ubiquitin-related modifier (SUMO) proteins, which include SUMO-1, SUMO-2 and SUMO-3, belong to the ubiquitin-like protein family. Like ubiquitin, the SUMO proteins are synthesized as precursor proteins that undergo processing before conjugation to target proteins. Also, both utilize the E1, E2, and E3 cascade enzymes for conjugation. However, SUMO and ubiquitin differ with respect to targeting. Ubiquitination predominantly targets proteins for degradation, whereas sumoylation targets proteins to a variety of cellular processing, including nuclear transport, transcriptional regulation, apoptosis and protein stability. The unconjugated SUMO-1, SUMO-2 and SUMO-3 proteins localize to the nuclear membrane, nuclear bodies and cytoplasm, respectively. SUMO-1 utilizes Ubc9 for conjugation to several target proteins, which include IkBa, MDM2, p53, PML and Ran GAP1. SUMO-2 and SUMO-3 contribute to a greater percentage of protein modification than does SUMO-1, and unlike SUMO-1, they can form polymeric chains. In addition, SUMO-3 regulates b-Amyloid generation and may be critical in the onset or progression of Alzheimera?s disease.

Note: This product is for in vitro research use only