

Myosin VIIa Antibody FITC Conjugated

Catalog No: #C07002F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Myosin VIIa Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 875-910 2215 derived from human Myosin VIIa
Conjugates	FITC
Target Name	Myosin VIIa
Other Names	DFNB2; MYU7A; NSRD2; USH1B; DFNA11; MYOVIIA; Unconventional myosin-VIIa; MYO7A
Accession No.	Swiss-Prot#Q13402NCBI Gene ID4647
Uniprot	Q13402
GeneID	4647;
Excitation Emission	494nm 518nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

IF=1:50-200

Background

Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails bind to membranous compartments, which are then moved relative to actin filaments. In the retina, plays an important role in the renewal of the outer photoreceptor disks. Plays an important role in the distribution and migration of retinal pigment epithelial (RPE) melanosomes and phagosomes, and in the regulation of opsin transport in retinal photoreceptors. In the inner ear, plays an important role in differentiation, morphogenesis and organization of cochlear hair cell bundles. Involved in hair-cell vesicle trafficking of aminoglycosides, which are known to induce ototoxicity (By similarity). Motor protein that is a part of the functional network formed by USH1C, USH1G, CDH23 and MYO7A that mediates mechanotransduction in cochlear hair cells. Required for normal hearing.

Note: This product is for in vitro research use only