Claudin 7 (Phospho-Tyr210) Antibody

Catalog No: #11794

Package Size: #11794-1 50ul #11794-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	Claudin 7 (Phospho-Tyr210) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of Claudin 7 only when phosphorylated at tyrosine 210.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 210 (S-K-E-Y(p)-V) derived from Human Claudin 7
Target Name	Claudin 7
Modification	Phospho
Other Names	CEPTRL2; claudin 7; CLD7; CPETRL2; CLDN7
Accession No.	Swiss-Prot#: O95471; NCBI Gene#: 1366; NCBI Protein#: NP_001171951.1.
Uniprot	O95471
GeneID	1366;
SDS-PAGE MW	32kd
Concentration	1.0mg/ml


Application Details

Western blotting: 1:500~1:1000

Images

Formulation

Storage



and 50% glycerol.

Store at -20°C/1 year

Western blot analysis of extracts from 3T3 cells (Lane 2), using Claudin 7 (Phospho-Tyr210) Antibody #11794. The lane on the left is treated with antigen-specific peptide.

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide

Background

Claudins, such as CLDN7, are involved in the formation of tight junctions between epithelial cells. Tight junctions restrict lateral diffusion of lipids and membrane proteins, and thereby physically define the border between the apical and basolateral compartments of epithelial cells.

Keen T.J.; Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases.

Kalnine N., Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

The MGC Project Team; Genome Res. 14:2121-2127(2004).

Note: This product is for in vitro research use only