MuSK (Phospho-Tyr755) Antibody

Catalog No: #11837

Package Size: #11837-1 50ul #11837-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description			
Product Name	MuSK (Phospho-Tyr755) Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.		
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho		
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.		
Applications	WB		
Species Reactivity	Hu		
Specificity	The antibody detects endogenous levels of MuSK only when phosphorylated at tyrosine 755.		
Immunogen Type	Peptide-KLH		
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 755(A-D-Y(p)-Y-K) derived from Human MuSK .		
Target Name	MuSK		
Modification	Phospho		
Other Names	MUSK; Muscle; skeletal;		
Accession No.	Swiss-Prot#: O15146; NCBI Gene#: 4593; NCBI Protein#: NP_005583.1.		
Uniprot	O15146		
GeneID	4593;		
SDS-PAGE MW	97kd		
Concentration	1.0mg/ml		
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide		
	and 50% glycerol.		
Storage	Store at -20°C/1 year		

Application Details

Western blotting: 1:500~1:1000

Images

250-	1 2	
130- 95- 72-	-	– MuSK (Phospho-Tyr755)
55-		
36- 28-		
17- 10- (Kd)		

Western blot analysis of extracts from JK cells (Lane 2), using MuSK (Phospho-Tyr755) Antibody #11837. The lane on the left is treated with antigen-specific peptide.

Background

Receptor tyrosine kinase which plays a central role in the formation and the maintenance of the neuromuscular junction (NMJ), the synapse between the motor neuron and the skeletal muscle. Recruitment of AGRIN by LRP4 to the MUSK signaling complex induces phosphorylation and activation of MUSK, the kinase of the complex. The activation of MUSK in myotubes regulates the formation of NMJs through the regulation of different processes including the specific expression of genes in subsynaptic nuclei, the reorganization of the actin cytoskeleton and the clustering of the acetylcholine receptors (AChR) in the postsynaptic membrane.

Valenzuela D.M., Neuron 15:573-584(1995).

Humphray S.J., Nature 429:369-374(2004).

Bergamin E., Mol. Cell 39:100-109(2010).

Note: This product is for in vitro research use only