

Arginase 1 Antibody FITC Conjugated

Catalog No: #C07578F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Arginase 1 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 184-220 322 derived from human Arginase 1
Conjugates	FITC
Target Name	Arginase 1
Other Names	Arginase-1; Liver-type arginase; Type I arginase; ARG1
Accession No.	Swiss-Prot#P05089NCBI Gene ID383
Uniprot	P05089
GeneID	383;
Excitation Emission	494nm 518nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1:50-200 ICC=1:50-200 IF=1:50-200

Background

Arginase I which is expressed almost exclusively in the liver, catalyzes the conversion of arginine to ornithine and urea. The human arginase I gene, which maps to chromosome 6q23, encodes a 322 amino acid protein. Arginase I exists as a homotrimeric protein and contains a binuclear manganese cluster. Arginase II catalyzes the same reaction as arginase I, but differs in its tissue specificity and subcellular location. Specifically, arginase II localizes to the mitochondria. Arginase II is expressed in non-hepatic tissues, with the highest levels of expression in the kidneys, but, unlike arginase I, is not expressed in liver. The human arginase II gene, which maps to chromosome 14q24.1-q24.3, encodes a 354 amino acid protein. In addition, arginase II contains a putative amino-terminal mitochondrial localization sequence.

Note: This product is for in vitro research use only