FXN Antibody

Catalog No: #32413

Package Size: #32413-1 50ul #32413-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| 00 | ori | nt | IOP | |
|----|-----|----|-----|--|

| Description | | |
|-----------------------|--|--|
| Product Name | FXN Antibody | |
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Purification | Antibodies were purified by affinity purification using immunogen. | |
| Applications | WB,IHC,IF | |
| Species Reactivity | Human,Mouse,Rat | |
| Specificity | The antibody detects endogenous level of total FXN protein. | |
| Immunogen Type | Recombinant Protein | |
| Immunogen Description | Recombinant protein of human FXN. | |
| Target Name | FXN | |
| Other Names | CyaY; FA; FARR; FRDA; MGC57199 | |
| Accession No. | Swiss-Prot:Q16595NCBI Gene ID:2395 | |
| Uniprot | Q16595 | |
| GeneID | 2395; | |
| SDS-PAGE MW | 23KD | |
| Concentration | 1.0mg/ml | |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% | |
| | sodium azide and 50% glycerol. | |
| Storage | Store at -20°C | |
| | | |


Application Details

WB 1:500 - 1:2000IHC 1:50 - 1:200IF 1:50 - 1:200


Images




Immunofluorescence analysis of C6 cells using FXN / Frataxin at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of L929 cells using FXN / Frataxin at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U-2 OS cells using FXN / Frataxin at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunohistochemistry of paraffin-embedded human mammary cancer using FXN / Frataxin at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human colon using FXN / Frataxin at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human lung cancer using FXN / Frataxin at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded rat lung using FXN / Frataxin at dilution of 1:100 (40x lens).

Immunohistochemistry of paraffin-embedded rat pancreas using FXN / Frataxin at dilution of 1:100 (40x lens).

Background

FXN, also named as FRDA, X25, m81-FXN, d-FXN, m78-FXN and i-FXN, belongs to the frataxin family. It promotes the biosynthesis of heme and assembly and repair of iron-sulfur clusters by delivering Fe2+ to proteins involved in these pathways. FXN may play a role in the protection against iron-catalyzed oxidative stress through its ability to catalyze the oxidation of Fe2+ to Fe3+; the oligomeric form but not the monomeric form has in vitro ferroxidase activity. FXN is cleaved to be 4 chains. The mature form of FXN is 14kd or 18kd.

Note: This product is for in vitro research use only