syk(Ab-323) Antibody

Catalog No: #21546

Package Size: #21546-1 50ul #21546-2 100ul





Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description           |   |  |  |  |
|-----------------------|---|--|--|--|
| Product Name          | syk(Ab-323) Antibody  |  |  |  |
| Host Species          | Rabbit  |  |  |  |
| Clonality             | Polyclonal  |  |  |  |
| Purification          | Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were |  |  |  |
|                       | purified by affinity-chromatography using epitope-specific peptide.                                       |  |  |  |
| Applications          | WB IHC IF   |  |  |  |
| Species Reactivity    | Hu  |  |  |  |
| Specificity           | The antibody detects endogenous level of total syk protein.   |  |  |  |
| Immunogen Type        | Peptide-KLH   |  |  |  |
| Immunogen Description | Peptide sequence around aa. 321~325 (N-P-Y-E-P) derived from Human syk.                                   |  |  |  |
| Target Name           | syk   |  |  |  |
| Other Names           | Spleen tyrosine kinase  |  |  |  |
| Accession No.         | Swiss-Prot: P43405NCBI Protein: NP_001128524.1  |  |  |  |
| Uniprot               | P43405  |  |  |  |
| GeneID                | 6850;   |  |  |  |
| Concentration         | 1.0mg/ml  |  |  |  |
| Formulation           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%      |  |  |  |
|                       | sodium azide and 50% glycerol.  |  |  |  |
| Storage               | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.                 |  |  |  |
|                       |   |  |  |  |


| Application Details            |    |  |  |
|--------------------------------|----|--|--|
| Predicted MW: 72kd             |    |  |  |
| Western blotting: 1:500~1:1000 |    |  |  |
| Immunohistochemistry: 1:50~1:  | 00 |  |  |
| Immunofluorescence: 1:100~1:2  | 00 |  |  |

## Images





Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using syk(Ab-323) Antibody #21546(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using syk(Ab-323) Antibody #21546.

## Background

Positive effector of BCR-stimulated responses. Couples the B-cell antigen receptor (BCR) to the mobilization of calcium ion either through a phosphoinositide 3-kinase-dependent pathway, when not phosphorylated on tyrosines of the linker region, or through a phospholipase C-gamma-dependent pathway, when phosphorylated on Tyr-348 and Tyr-352. Thus the differential phosphorylation of Syk can determine the pathway by which BCR is coupled to the regulation of intracellular calcium ion

Zhang, J. et al. (2000) J. Biol. Chem. 275, 35442-35447.

Turner, M. et al. (2000) Immunol. Today 21, 148-154.

Decker, M. et al. (1998) J. Biol. Chem. 273, 8867-8874.

Law, C.L. et al. (1996) Mol. Cell. Biol. 16, 1305-1315.

Note: This product is for in vitro research use only