

BRCA1(Phospho-Ser988) Antibody

Catalog No: #11237

Package Size: #11237-1 50ul #11237-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


Product Name	BRCA1(Phospho-Ser988) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	IF
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of BRCA1 only when phosphorylated at serine 988.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 988 (I-K-S(p)-F-V) derived from Human BRCA1.
Target Name	BRCA1
Modification	Phospho
Other Names	RNF53
Accession No.	Swiss-Prot: P38398NCBI Protein: NP_009225.1
Uniprot	P38398
GeneID	672;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 220kd

Immunofluorescence: 1:100~1:200

Images



Immunofluorescence staining of methanol-fixed HeLa cells using BRCA1(Phospho-Ser988) Antibody #11237.

Background

The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. Plays a central role in DNA repair by facilitating cellular response to DNA repair. Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. Involved in transcriptional regulation of P21 in response to DNA damage. Required for FANCD2 targeting to sites of DNA damage. May function as a transcriptional regulator. Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation

Lin SY, et al. (2004) Proc Natl Acad Sci U S A, 101(17): 6484-6489

Zhang J, et al. (2004) Mol Cell Biol; 24(2): 708-718

Beger C, et al. (2001) Proc Natl Acad Sci U S A; 98(1): 130-135

Gardner K, et al. (2001) Breast Cancer Res; 3(1): 11-13

Zheng L, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9587-9592

Note: This product is for in vitro research use only