Adiponectin Antibody

Catalog No: #21665

Package Size: #21665-1 50ul #21665-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$			
	escri	ntı	Λn
$\boldsymbol{ u}$	COUL	рu	OH


Product Name	Adiponectin Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB	
Species Reactivity	Hu Ms Rt	
Specificity	The antibody detects endogenous level of total Adiponectin protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa.235~239(T-G-F-L-L) derived from Human Adiponectin.	
Target Name	Adiponectin	
Other Names	ACDC; ADPN; APM1	
Accession No.	Swiss-Prot: Q15848NCBI Protein: NP_001171271.1	
Uniprot	Q15848	
GeneID	9370;	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

Application Details

Predicted MW: 27kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from Rat and Mouse brain tissue using Adiponectin Antibody #21665.

Background

Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-a by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW. Yamauchi T., Kamon J., Waki H. Kadowaki T.Nat. Med. 7:941-946(2001)

Richards A.A., Stephens T., Charlton H.K. Mol. Endocrinol. 20:1673-1687(2006)

Note: This product is for in vitro research use only