USP2 Antibody

Catalog No: #36169



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_			
\mathbf{I}	esci	unt	ion.
$\boldsymbol{-}$	しつし	IΙDι	IUII

finity purification.
dy detects endogenous levels of total USP2 protein.
ant Protein
tein corresponding to residues near the C terminal of human ubiquitin specific peptidase 2
P41
#: O75604NCBI Gene ID: 9099Gene Accssion: BC002854
in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
0°C
, i


Application Details

Immunohistochemistry: 1:50-1:200

Images



Immunohistochemical analysis of paraffin-embedded Human cervical cancer tissue using #36169 at dilution 1/40.



Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using #36169 at dilution 1/40.

Background

This gene encodes a member of the family of de-ubiquitinating enzymes, which belongs to the peptidase C19 superfamily. The encoded protein is a ubiquitin-specific protease which is required for TNF-alpha (tumor necrosis factor alpha) -induced NF-kB (nuclear factor kB) signaling. This protein deubiquitinates polyubiquitinated target proteins such as fatty acid synthase, murine double minute 2 (MDM2), MDM4/MDMX and cyclin D1. MDM2 and MDM4 are negative regulators of the p53 tumor suppressor and cyclin D1 is required for cell cycle G1/S transition. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Note: This product is for in vitro research use only